## **NEW YORK STATE** OFFICE OF CHILDREN AND FAMILY SERVICES INDIVIDUAL ALLERGY AND ANAPHYLAXIS EMERGENCY PLAN

## Instructions:

- This form is to be completed for any child with a known allergy.
- The child care program must work with the parent(s)/guardian(s) and the child's health care provider to develop written instructions outlining what the child is allergic to and the prevention strategies and steps that must be taken if the child is exposed to a known allergen or is showing symptoms of exposure.
- This plan must be reviewed upon admission, annually thereafter, and anytime there are staff or volunteer changes, and/or anytime information regarding the child's allergy or treatment changes. This document must be attached to the child's Individual Health Care Plan.
- Add additional sheets if additional documentation or instruction is n

Asthma: ☐ Yes (h <b>Wy child is reactive</b>	Current Weight: igher risk for reaction) \( \subseteq \text{No} \) to the following allergens:	lbs.
Allergen:	Type of Exposure: (i.e., air/skin contact/ingestion, etc.):	Symptoms include but are not limited to: (check all that apply)
		☐ Shortness of breath, wheezing, or coughing ☐ Pale or bluish skin, faintness, weak pulse, dizziness. ☐ Tight or hoarse throat, trouble breathing or swallowing ☐ Significant swelling of the tongue or lips ☐ Many hives over the body, widespread redness ☐ Vomiting, diarrhea ☐ Behavioral changes and inconsolable crying
		☐ Shortness of breath, wheezing, or coughing ☐ Pale or bluish skin, faintness, weak pulse, dizziness ☐ Tight or hoarse throat, trouble breathing or swallowing
		☐ Significant swelling of the tongue or lips ☐ Many hives over the body, widespread redness ☐ Vomiting, diarrhea ☐ Behavioral changes and inconsolable crying ☐ Other (specify)
		☐ Shortness of breath, wheezing, or coughing ☐ Pale or bluish skin, faintness, weak pulse, dizziness ☐ Tight or hoarse throat, trouble breathing or swallowing ☐ Significant swelling of the tongue or lips ☐ Many hives over the body, widespread redness ☐ Vomiting, diarrhea
ny child was LIKEL)	l i	Behavioral changes and inconsolable crying
ny child was DEFINI	TELY exposed to an allorgon are seen	
☐ give epinephrin	e immediately	- 3broup are bresent:

OCFS-6029 (12/2020)	
Date of Plan; / /	'C
THE FOLLOWING STEPS WILL BE TAKEN IF THE CHILD EXHIBITS SYMPTOMS including, but not limited to:  On the state of the state	
Inject eninenhring insertition	
<ul> <li>Inject epinephrine immediately and note the time when the first dose is given.</li> <li>Call 911/local rescue squad (Advise 911 the child is in anaphylaxis and may need epinephrine when emerged and the person flet and the person</li></ul>	
responders arrive).	enci
or lie on their side.	
<ul> <li>If symptoms do not improve, or symptoms return, an additional dose of epinephrine can be given in consultation.</li> </ul>	up
Alert the child's parents/guardians and emergency contacts.      After the poods of the child.	on
• After the needs of the child and all all all and all all all all all all all all all al	
After the needs of the child and all others in care have been met, immediately notify the office.  MEDICATION/DOCSES	
WED10/4/10/6/00252	
Epinephrine brand or generic:	
Epinephrine dose: □ 0.1 mg IM □ 0.15 mg IM □ 0.3 mg IM	
ADMINISTRATION AND SAFETY INFORMATION FOR EPINEPHRINE AUTO-INJECTORS  When administering an epinephrine auto-injector follows to a construction of the construction of	
When administering an epinephrine auto-injector follow these guidelines:	
<ul> <li>Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than the emergency room.</li> <li>If administration are the statement of the statemen</li></ul>	ne
If administering an auto-injector to a young child, hold their leg firmly in place before and during injection to	
prevent injuries.	
Epineprinne can be injected through clothing if pand at	
Call 911 immediately after injection.	
STORAGE OF EPINEPHRINE AUTO-INJECTORS	
All medication will be be a six to	
All medication will be kept in its original labeled container.      Medication must be kept in its original labeled container.	
meandation must be kept in a clean area that is increased.	
The state of the s	
modifications, such as enthernrine outs initiative	
<ul> <li>Explain here where medication will be stored:</li> </ul>	
MAT/EMAT CERTIFIED PROGRAMS ONLY	
Oliv Sidil listed in the near the second side of th	
medications. Staff must be at least 18 years old and have first aid and ODD	
and of Acel Inicales that cover all agon of ability	
drand of generic:	
Antihistamine dose:	
Other (e.g., inhaler-bronchodilator if wheezing):	
*Note: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE	
STORAGE OF INHALERS ANTIHISTAMMED PROCESS	
All medication will be kept in its original label.	
All medication will be kept in its original labeled container. Medication must be kept in a clean area that is inaccessible to stored. Note any medications, such as as three into the child's medication is stored. Explain where	
children. All staff must have an awareness of where the child's medication is stored. Note any medications, such as asthma inhalers, that may be stored in a different area.  Explain here:	

STRATEGIES TO REDUCE THE RISK OF EXPOSURE TO ALLERGIC TRIGGERS

The following strategies will be taken by the child care program to minimize the risk of exposure to any allergens while the above-named child is in care (add additional sheets if needed):

Document plan here:	aou).	
EMERGENCY CONTACTS - CALL 911		
Ambulance: ( ) -		
Child's Health Care Provider:		
Parent/Guardian:	Phone #: (	) -
CHILD'S EMERGENCY CONTACTS	Phone #: (	) -
Name/Relationship:		
Name/Relationship:	Phone#: (	) -
lame/Relationship:	Phone#: (	
or conditionally.	Phone#: (	) -
arent/Guardian Authorization Signature:	. Horien. (	
hysician/HCD Authority:	Date:	
hysician/HCP Authorization Signature:		
rogram Authorization Signature:	Date:	
	Date:	_ / /